

Department of Economics

University of Manitoba

ECON 7010: Econometrics I
FINAL EXAM, Dec. 12th, 2020

Instructor: Ryan Godwin
Instructions: Upload your answers to UM Learn.
Time Allowed: 2 hours.
Number of Pages: 3

There are a total of 100 marks.

Answer all questions. Each question is worth 20 marks.

1.) The assumptions underlying the classical linear regression model are:

- A.1 Linearity
- A.2 Full rank: $\text{rank}(X) = k$
- A.3 Errors have zero mean: $E(\varepsilon) = 0$
- A.4 Spherical errors
- A.5 The process that generates X is unrelated to the process that generates ε
- A.6 Normality of errors

Under these assumptions, OLS is a good estimator. Additionally, $s^2(X'X)^{-1}$ is a good estimator for the variance-covariance matrix of the OLS estimator. For each of the assumptions listed above (if applicable): (i) describe the consequences of the assumption being violated (in terms of estimation, hypothesis testing, etc.), and (ii) describe how the problem might be corrected, or how an alternative estimator might be used to correct the problem (if possible).

2.) Suppose that we have a linear multiple regression model,

$$y = X\beta + \varepsilon,$$

where the regressors may be random and may be correlated with the errors, even asymptotically. Accordingly, we decide to use a (generalized) IV estimator, where Z is the $(n \times L)$ matrix of (possibly random) instruments.

a) Prove, algebraically, that the IV estimator is equivalent to the following two-step estimator:

- (i) Regress X on Z by OLS, and get the predicted matrix, \hat{X} .
- (ii) Fit the following artificial model by OLS: $y = \hat{X}\beta + v$.

b) Prove that the generalized IV estimator collapses to the simple one, if X and Z have the same dimensions.

c) Briefly describe how you would test to see if IV estimation is needed.

d) Is there a way to test if the assumption that $\text{plim}(Z'\varepsilon/n) = 0$ is satisfied? Explain.

3.) Consider following wage equation:

$$\log(\text{wage}) = \beta_0 + \beta_1 \text{Educ} + \beta_2 \text{Exper} + \beta_3 \text{Marr} + \epsilon$$

where wage is hourly wage, Educ is years of education, Exper is job experience in years, and Marr is a dummy variable that is equal to 1 if individual is married and zero if single.

a) Using the estimation results in the table below calculate the **F-statistics** in order to test the following hypothesis:

$$H_0: \beta_3 = 0.5$$

Explain your reasoning.

```
##
## =====
##                               Dependent variable:
##                               -----
##                               lwage
## -----
## educ                          0.09***
##                               (0.01)
##
## exper                          0.01***
##                               (0.002)
##
## married                        0.19***
##                               (0.04)
##
## Constant                       0.22**
##                               (0.11)
##
## -----
## Observations                   526
## R2                             0.28
## Adjusted R2                    0.27
## Residual Std. Error            0.45 (df = 522)
## F Statistic                    66.27*** (df = 3; 522)
## =====
## Note:                          *p<0.1; **p<0.05; ***p<0.01
##                               Standard Errors are in parenthesis
```

b) Assuming that you cannot reject the following hypothesis

$$H_0: \beta_1 - 9\beta_2 = 0, \text{ and } \beta_3 = 1$$

Derive the restricted model. Why is the LS estimator of the restricted model more appropriate than the unrestricted model? Explain.

c) As we learned in class the hypothesis given in section (b) can be re-written as

$$H_0: R\beta = q$$

We also saw that $m \sim N(0, V(m))$ where $m = Rb - q$

Calculate the **Wald Statistic** that can be used to test the hypothesis in section (b) assuming that a consistent estimate of $V(m)^{-1}$ is given below.

```
5443  -481
-481   572
```

4.) Suppose that $V(\boldsymbol{\varepsilon}) \neq \sigma^2 I$. Instead, heteroskedasticity is present and $V(\boldsymbol{\varepsilon}) = \sigma^2 \Omega$.

a) Explain the implications of ignoring heteroskedasticity.

b) Derive $V(\mathbf{b})$ for when $V(\boldsymbol{\varepsilon}) = \sigma^2 \Omega$.

c) Describe White's test for heteroskedasticity.

d) Suppose that you have an additional variable, z_i , which describes the form of heteroskedasticity, such that:

$$\Omega = \begin{bmatrix} z_1 & 0 & 0 & \dots & 0 \\ 0 & z_2 & 0 & \dots & 0 \\ 0 & 0 & z_3 & \dots & 0 \\ \vdots & \vdots & & \ddots & \vdots \\ 0 & 0 & \dots & & z_n \end{bmatrix}$$

Explain how to transform the model into one that is characteristic of homoskedasticity, and how you could obtain the Generalized Least Squares (GLS) estimates from the transformed model.

e) Prove that the transformed model from (d) above satisfies A.4.

5.)

a) Suppose that we wish to estimate a regression model by least squares, but that this model is non-linear in the parameters. Will we need to use a numerical algorithm to obtain the parameter estimates? If so, explain why.

b) Illustrate how the Newton-Raphson algorithm can find the value of θ that minimizes the function:

$$f(\theta) = \frac{\theta^4}{2} - \frac{\theta^3}{3}$$

END.